



ΕΛΛΗΝΙΚΗ ΜΑΙΕΥΤΙΚΗ & ΓΥΝΑΙΚΟΛΟΓΙΑ

www.iatrikionline.gr




TABLE III
DEATH-RATE FOR ALL BLOOD-POSITIVE CASES (HÆMOLYTIC STREPTOCOCCI) WITH OR WITHOUT GENERALISED PERITONITIS

Year.	Cases.	Deaths.	Death-rate (per cent.).
1932	14	13	93
1933	24	18	75
1934	26	15	58
1935	18	12	67
Totals	82	58	Aver. 71
1936 and first* quarter of 1937 (Prontosil and sulph-anilamide treatment) ..	22	6	27.3

* The cases for 1936 and first quarter of 1937 only were chosen because Gibberd's analysis (1937) had dealt with that period.

ΠΑΡΑΡΤΗΜΕΝΟ
ΤΕΛΟΣ
ΤΟΥ ΓΡΑΦΕΙΟΥ
ΚΕΜΠΤΑ
Αρ. Πρωτ. 794/94



ΜΕΣΟΓΕΙΩΝ 215 115 25 ΑΘΗΝΑ

HELLENIC OBSTETRICS AND GYNECOLOGY

ΤΟΜΟΣ 22 - ΤΕΥΧΟΣ 4 - 2010

VOLUME 22 - No 4 - 2010

ΠΕΡΙΟΔΙΚΗ ΤΡΙΜΗΝΙΑΙΑ ΕΚΔΟΣΗ - ΕΠΙΣΗΜΟ ΟΡΓΑΝΟ ΤΗΣ
- ΜΑΙΕΥΤΙΚΗΣ & ΓΥΝΑΙΚΟΛΟΓΙΚΗΣ ΕΤΑΙΡΕΙΑΣ ΘΕΣΣΑΛΟΝΙΚΗΣ
- ΜΑΙΕΥΤΙΚΗΣ & ΓΥΝΑΙΚΟΛΟΓΙΚΗΣ ΕΤΑΙΡΕΙΑΣ ΚΡΗΤΗΣ
- ΜΑΙΕΥΤΙΚΗΣ & ΓΥΝΑΙΚΟΛΟΓΙΚΗΣ ΕΤΑΙΡΕΙΑΣ ΚΕΝΤΡΙΚΗΣ ΕΛΛΑΔΑΣ
- ΜΑΙΕΥΤΙΚΗΣ & ΓΥΝΑΙΚΟΛΟΓΙΚΗΣ ΕΤΑΙΡΕΙΑΣ ΑΝΑΤ. ΜΑΚΕΔΟΝΙΑΣ & ΘΡΑΚΗΣ
- ΕΛΛΗΝΙΚΗΣ ΕΤΑΙΡΕΙΑΣ ΕΝΔΟΜΗΤΡΙΩΣΗΣ

PUBLISHED QUARTERLY - OFFICIAL JOURNAL OF THE
- OBSTETRICAL AND GYNECOLOGICAL SOCIETY OF THESSALONIKI
- OBSTETRICAL AND GYNECOLOGICAL SOCIETY OF CRETE
- OBSTETRICAL AND GYNECOLOGICAL SOCIETY OF CENTRAL GREECE
- OBSTETRICAL AND GYNECOLOGICAL SOCIETY OF EASTERN MACEDONIA & THRACE
- HELLENIC SOCIETY OF ENDOMETRIOSIS